

Specialists In Reproductive Medicine & Surgery, P.A.

www.DreamABaby.com • Fertility@DreamABaby.com

Excellence, Experience & Ethics



Early Pregnancy Patient Instructions

Congratulations on your pregnancy! Below are some guidelines to assist you in some of the decision-making at this important time. These are only basic recommendations so please contact us if you have more specific questions.

Obstetrical Ultrasounds:

- We usually recommend an ultrasound at approximately 4 weeks after the time of *insemination* or probable *conception*, or 6 weeks from the first day of your last menstrual period. The ultrasound is frequently repeated again two weeks later.
- If you were provided with any videos during the previous procedures, please bring these all cued up to the end of the last recording so that we can continue to record your ultrasound on your videotape. ***Please remember to bring your tape to every ultrasound visit!***
- If you so desire, you may tell close family members about your pregnancy after your first ultrasound, but we usually suggest that you wait to tell other friends and family until your second ultrasound confirms continued embryonic/fetal growth.

Bleeding:

- The early embryo (and often, ***EMBRYOS!***) burrow into the uterine lining and cause implantation bleeding. A miscarriage, however, involves significant cramping and bleeding (equal to and often greater than a heavy period) and is quite different from implantation bleeding. ***Please do not call the physician during off hours for implantation bleeding.*** Implantation bleeding is not an emergency and is very, very common. In order to follow you appropriately, however, please do notify the office during regular business hours that implantation bleeding occurred.
- Later in the pregnancy (> 6 weeks from the last menstrual period), when the embryo/fetus has created their own red blood cells, vaginal bleeding can lead to the mixing of the maternal and fetal blood. If the fetus is Rh+ and the mother Rh-, an injection of RhoGAM will need to be administered to the mother within three days of the initial bleed. Please contact the nurses on the first business day should bleeding occur after six weeks gestational age.
- If significant bleeding occurs with cramping, you may indeed be trying to miscarry. Please contact the physician on call when this occurs. Understand, however, little if anything can be done to change the course of events. All that can be done is to make sure that you are safe and comfortable during the spontaneous loss of the pregnancy or during a scheduled evacuation of the uterine contents.

Medications:

- Begin taking prenatal vitamins immediately if you are not already doing so. Please ask us if you do not have a prescription for these. Taking your prenatal vitamin with food or before bedtime with a snack will minimize stomach upset.
- Discontinue all medications except your pre-natal vitamins and those specifically prescribed by a physician. (Make sure the physician prescribing knows you are pregnant.)
- If you are taking Progesterone, ask us about your specific instructions.
- For pain relief, it is safe to take **Extra-Strength Tylenol** as directed on the package. Do not take **Advil**, **Aspirin**, or other pain remedies unless specifically instructed by your physician.
- For mild cold symptoms, **Tylenol**, plain **Robitussin**, **Chlor-Trimeton** or **Benadryl** may be taken. If you have a thyroid problem, please do not take the **Robitussin** without permission. **Chloraseptic Spray/Lozenges**, **Cepastat Lozenges** and cough drops are also allowed. **Afrin Nasal Spray** may be used for a very short duration for nasal congestion. It is best, however, to avoid all medications during the first 12 weeks of gestation except those specifically prescribed by your physician or any physician who is aware of your pregnancy. Any medications prescribed by a non-OB/GYN should be cleared by your OB/GYN before taking and we encourage your physician to call Specialists in Reproductive Medicine & Surgery, P.A., (SRMS) if they have any particular questions.

Diet: Nausea/Vomiting:

Nausea and some vomiting during the first trimester are quite common. If you have these problems, please try some of the remedies listed below:

- If you are unable to keep any fluids down over a twenty-four hour period or show early signs of dehydration, such as decreased urination, please call the office.
- Avoid spicy, greasy, fatty, heavy or rich foods.
- Eating a cracker before rising has been helpful to some women with morning nausea.
- Eat smaller and more frequent meals. Avoid large meals.
- Drink lots of fluids but avoid caffeinated fluids since caffeine may be associated with increased miscarriages.
- Avoid smells that might trigger nausea such as smoke and smelly foods.
- Marine seasickness wristbands have helped some pregnant women control their nausea. These are available at some pharmacies and boating stores.
- Take your prenatal vitamin with food or before bedtime with a snack. If you are unable to take your vitamin, please try to take two chewable vitamins each day. Please make certain that you are consuming at least 400 mcg. (0.4 mg) of Folate (also called folic acid) each day.
- If you are having indigestion, you are free to take **Tums**, **Riopan**, **Riopan Plus** and **Mylanta** as directed on the package. Pregnancy-associated indigestion may also be improved with the use of chewing gum for 30 minutes following meals and as needed.
- There is data that indicates that substances such as alcohol, smoking and caffeine increase your chances for miscarriage and fetal malformations. It is best to minimize your exposure to the above substances. There are really no known safe levels of any of these substances.

Diet: Constipation

- Constipation may occur frequently during your pregnancy.
- Eat a high fiber diet.
- Some women suggest prune juice (easier to tolerate taste if warmed), green apples, or fruit juices.

Early Pregnancy Patient Instructions (*cont.*)

- Over the counter fiber remedies may be used when the above do not help. These include *Colace* 200 mg twice each day, *Metamucil*, *Citricil* or *Milk of Magnesia* 2 tsp. at bedtime.
- Call the office for severe constipation.

Activity/Exercise:

- You may continue activities that you were involved in prior to the pregnancy.
- It is best to not to start a new exercise program.
- Do not horse back ride, scuba dive or involve yourself in other high-risk sports.
- Walking and swimming are great forms of exercise during pregnancy.
- Be sure to drink plenty of fluids to prevent dehydration from developing.

Breast Tenderness:

Breast tenderness is often a welcome sign of pregnancy. Unfortunately, once present, there are a limited number of ways to minimize the discomfort.

- Wear a supportive bra.
- Take *Extra-Strength Tylenol* according to package instructions.

Sexual Intercourse:

- Vaginal intercourse is allowed if you are not actively bleeding.
- If you have lost previous pregnancies or are unusually anxious about your current pregnancy, SRMS suggests that the patients decide if vaginal intercourse is allowed. Vaginal penetration is not thought to cause a spontaneous abortion, but the emotional consequences of intercourse, followed by spontaneous bleeding and loss can be difficult. It is generally suggested that couples engage in sexual activities if they feel comfortable with the fact that the activity will, in and of itself, not cause a spontaneous loss.

History of Hypothyroidism:

Under-treated hypothyroidism can occur in pregnant women because of the rapid needs of the fetus for thyroid hormone. Increased requirements for thyroid hormone occur as early as the fifth week of pregnancy. Under-treated hypothyroidism has been shown to triple a woman's chance of placental abruption (the placenta pulling away from the uterine wall) and double the risk of preterm birth. Infants of women with under-treated hypothyroid have twice the risk of respiratory distress and admission to a neonatal intensive care unit. For these reasons, the following will be recommended:

- With a positive Quantitative hCG (blood pregnancy test), we will advise you to take an extra dose of your thyroid hormone on Tuesday and Friday of each week.
- We will obtain a TSH and Free T4 level along with your second Quantitative HCG level.
- We will monitor your TSH and Free T4 levels every 4 weeks until 16-20 weeks of pregnancy.

Other Recommendations:

- Do not sit in hot tubs, hot baths, or saunas. Increasing a pregnant women's core body temperature to 102 degrees has been associated with embryonic/fetal malformations. It is okay to take warm showers, however.
- If you own or care for an outdoor cat, do not change the litter box. Cats may carry toxoplasmosis. If you want to see if you are already immune to toxoplasmosis, please ask your physician and we will be happy to obtain an antibody titer and determine immunity.
- Do not eat raw meat or fish during your pregnancy since there is an increased risk of parasites.
- Do not eat shark, swordfish, king mackerel, or tilefish during your pregnancy as they are high in mercury. You may eat up to 12 ounces or two average servings per week of fish low in mercury, such as salmon, shrimp, Pollack and catfish. Canned light tuna has less mercury than albacore. For local fish, check local advisories that apply to your area or limit ingestion to 6 ounces per week.
- Because of the small, but serious risk of listeriosis, a bacteria that can cause miscarriage and stillbirth, we recommend avoiding hot dogs or luncheon meats unless they have been heated to steaming. Also avoid soft cheeses, refrigerated pate's or meat spreads, and unpasteurized milk.
- If you smoke, quit! Try to avoid smoky atmospheres, as this has been associated with miscarriages.
- Do not scuba dive while pregnant.
- No association between hair dye and fetal malformations has been found, so this activity can be done if you desire.

Call The Office Immediately For:

- Fever (Temperature of 100.4 degrees Fahrenheit or greater, taken twice, four hours apart)
- Significant vaginal bleeding
- Abdominal pain
- Dizziness, severe headaches
- Persistent nausea and vomiting

We are so happy we could be a part of this miracle! Please accept our thanks for your trust in us. Remember to call the office and let us know how the delivery goes. We would also appreciate a picture for our success board and stories for the Internet are wonderful additions!

***Get plenty of rest!
Use good common sense!
Enjoy this very special time!***

**Thank you again for choosing
Specialists In Reproductive Medicine & Surgery, P.A!**